



## SPECIFICATION

## CHINESE HERBAL MEDICAL COMPOSITION IN THE FORM OF JELLY

## 5 TECHNICAL FIELD OF THE INVENTION

The present invention relates to a Chinese herbal medical composition in the form of jelly, which hardly causes syneresis, is superior in the preservative stability, is broadly applicable to Chinese herbal medicine (漢方藥), and can be orally taken without taking care of the bitter taste, etc., of the Chinese herbal medicine.

## BACKGROUND OF THE INVENTION

The traditional Chinese herbal medicines are in the forms of liquids prepared by decocting crude drugs (生藥), powders prepared by powdering crude drugs or pills made of crude drugs and honey. Therefore, traditional Chinese herbal medicines have the disadvantages of being inconvenient due to the time required to decoct crude drugs, and due to having to prepare it according to necessity. Furthermore, it is painful or difficult for a patient to take the decoction or the powdered crude drug, due to the bitter taste or smell, which are peculiar to Chinese herbal medicines. Today, in order to solve such problems, Chinese herbal medical preparations, such as extracts from a Chinese herbal medicine, powders, granules, tablets, liquids, etc., are prepared starting from the powdered crude drugs. These preparations solve the inconveniences due to taking time to decoct crude drugs, and having to prepare it according to necessity. These preparations are also superior in preservative stability.

However, a patient must take several grams of preparation in the

form of powders, granules or tablets, and this is a burden for the patient. Further, the powders and the granules can cause problems such as choking, resulting in a sandy feeling in a mouth, or getting between false teeth when they are taken. The tablets can be too large  
5 for a patient to take. The taste and smell peculiar to a Chinese herbal medicine is improved by tableting it. However, there are still disadvantages in that a Chinese herbal medicine is strongly tasted, is unpleasant, and is difficult to take when the contents in the tablets dissolve, or the tablets disintegrate in a mouth while taking them.

10 On the other hand, the liquids are more easily taken compared to the powders, the granules and the tablets. However, the liquids have troubles in that the bitter taste and smell peculiar to a Chinese herbal medicine becomes strong, because the liquid is broadly spread in a mouth. Therefore, it is painful and difficult to take a liquid formulation.  
15 Furthermore, it is inconvenient to carry as it is packed in a glass-bottle. In order to solve such problems with a Chinese herbal medical preparation, it is considered to make a Chinese herbal medicine in the form of jelly.

As a jelly preparation containing a Chinese herbal medicine, there  
20 is known a jelly preparation made of a Chinese herbal medicine and gelatin (Japanese patent publication B 7-116049). As gelatin is a gelling agent which is physico-chemically unstable, the preparation lacks preservative stability and must be stored in a cold place. Therefore, it does not stand for the test for medicines (the long term-  
25 preservation test at 25°C under 60 % RH for 3 years, or the accelerated preservation test at 40°C under 75 % RH for 6 months, etc.). In addition, gelatin is easily dissolved in a mouth and therefore, the preparation easily gives the bitter taste and is difficult to take when a

Chinese herbal medicine having the strong bitter taste is contained.

In addition, it is considered that a jelly preparation containing Chinese herbal medicine is prepared using sodium alginate or agar. As a jelly preparation containing a Chinese herbal medicine using sodium alginate, a jelly preparation containing Sho-saiko-to is known (小柴胡湯) (Japanese patent No.2508547). The bitter taste peculiar to a Chinese herbal medicine can be masked by adding alginic acid. However, as jelly containing alginic acid causes much syneresis, heterogeneity of the drug occurs and the drug remains in the packed vessel when taking it. The appearance is also bad. Therefore, the preparation is not preferable as a medicine.

In regard to a jelly preparation prepared using agar, the preparation causes much syneresis, similar to the preparation containing alginic acid. Further, the preparation easily disintegrates in a mouth, easily gives the bitter taste and does not give a good feeling when taking it.

As other jelly preparations, a jelly composition (Japanese patent publication A 9-187233 and Japanese patent publication A 9-194346) and a Chinese herbal medical composition in the form of jelly (Japanese patent publication A 2001-114696) are known, but it is very difficult to prepare jelly preparations containing a Chinese herbal medicine which guarantees the preservative stability on a medical level.

The following reasons are considered as causes which make it difficult to make a Chinese herbal medicine in the form of jelly.

The preparation contains as a starting material, a natural product which consists of a variety of ingredients, and many of these ingredients are structurally unknown. In addition, there are many forms of starting materials, such as crude drugs, liquid extract, condensed extract, dry

extract, soft extract, fluid extract, etc., and the dosages on them vary.

As there is such a background peculiar to a Chinese herbal medicine, when a Chinese herbal medicine is formed into a jelly preparation, according to the kind or amount of the contained Chinese herbal medicine, there are possibilities to produce the preparation wherein its appearance can not be maintained due to syneresis and the stability of the active ingredients can not be maintained. Therefore, a Chinese herbal medical composition in the form of jelly which is broadly applicable has been desired.

#### BRIEF SUMMARY OF THE INVENTION

The present invention was completed based on the above viewpoint and the goal is to provide a Chinese herbal medical composition in the form of jelly, which hardly causes syneresis, is superior in the preservative stability, is broadly applicable to a Chinese herbal medicine, and is orally taken without taking care of the bitter, etc., of a Chinese herbal medicine.

The present inventors have extensively studied to solve the above problems, and have found that by using at least one substance selected from the group consisting of carrageenan, carob bean gum and xanthan gum as a base (not containing phosphate buffer) of the jelly preparation containing a Chinese herbal medicine, a Chinese herbal medical composition in the form of jelly, which hardly causes syneresis, is superior in the preservative stability, is broadly applicable to a Chinese herbal medicine, and is orally taken without taking care of the bitter, etc., of a Chinese herbal medicine is obtainable. Thus, the present invention was completed.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a stick-like packed vessel which is sealed on three parts.

### Explanation of signs

- 5           1: A Chinese herbal medical composition in the form of jelly
- 2: Sealed parts

## DETAILED DESCRIPTION OF THE INVENTION

Carrageenan used in the Chinese herbal medical composition in the form of jelly of the present invention is not limited as long as it is usually used for a jelly composition. For example, there are  $\kappa$  (kappa) type,  $\iota$  (iota) type and  $\lambda$  (lambda) type, with respect to carrageenan. Any type is usable, but preferably a combination of  $\iota$  type carrageenan and either  $\kappa$  type carrageenan or  $\lambda$  type carrageenan, or a combination of these three type carrageenans, or  $\iota$  type carrageenan solely is usable. When a combination of  $\iota$  type carrageenan and either  $\kappa$  type carrageenan or  $\lambda$  type carrageenan, or a combination of these three type carrageenans is used,  $\iota$  type carrageenan is usually used in the amount of more than 50 w/w % per total carrageenan weight, preferably more than 70 w/w %, and more preferably more than 95 w/w %.

The amount of carrageenan contained in the Chinese herbal medical composition in the form of jelly of the present invention is preferably 0.01~10.0 w/w % per total amount of the composition, more preferably 0.05~5.0 w/w %, and further more preferably 0.08~2.0 w/w %. When the amount of carrageenan is beyond the above range, the preparation becomes too viscous to prepare it, and when the amount is below the above range, the jelly formation becomes difficult and the desired composition is not obtainable.

Carob bean gum used in the Chinese herbal medical composition in the form of jelly of the present invention is not limited as long as it is usually used for a jelly composition.

5 The amount of carob bean gum contained in the Chinese herbal medical composition in the form of jelly of the present invention is preferably 0.01~10.0 w/w % per total amount of the composition, more preferably 0.05~5.0 w/w %, and further more preferably 0.1~2.0 w/w %. When the amount of carob bean gum is beyond the above range, the preparation becomes too viscous to prepare it, and when the amount is  
10 below the above range, the jelly formation becomes difficult and the desired composition is not obtainable.

Xanthan gum used in the Chinese herbal medical composition in the form of jelly of the present invention is not limited as long as it is usually used for a jelly composition.

15 The amount of xanthan gum contained in the Chinese herbal medical composition in the form of jelly of the present invention is preferably 0.01~10.0 w/w % per total amount of the composition, more preferably 0.05~5.0 w/w %, and further more preferably 0.08~2.0 w/w %. When the amount of xanthan gum is beyond the above range,  
20 the preparation becomes too viscous to prepare it, and when the amount is below the above range, the jelly formation becomes difficult and the desired composition is not obtainable.

The amount of a Chinese herbal medicine contained in the Chinese herbal medical composition in the form of jelly of the present  
25 invention is preferably less than 60 w/w % per total amount of the composition, more preferably less than 50 w/w %, and further more preferably less than 30 w/w %. When the amount of the Chinese herbal medicine is beyond the above range, there is a possibility that

the preparation becomes too viscous to prepare it and the jelly formation becomes difficult due to it being rice cake or jam-like.

A dispersion medium in order to disperse a base which is used for a Chinese herbal medical composition in the form of jelly of the present invention includes a liquid which is usually used as a dispersion medium of a jelly composition, for example water or a mixture of water and a polyalcohol. Examples of the polyalcohol are glycerin, propylene glycol, etc. The amount of the dispersion medium of a Chinese herbal medical composition in the form of jelly of the present invention is, preferably 30~90 w/w % per total composition, more preferably 30~85 w/w %, and further more preferably 40~80 w/w %.

The Chinese herbal medical composition in the form of jelly of the present invention may, if necessary, contain various known ingredients which are acceptable as medical additives and are orally administrable, such as stabilizing agents, correctives, sweetening agents, emulsifying agents, dispersion agents, preservatives, flavors, coloring agents, etc.

The stabilizing agent, if desired, contained in the Chinese herbal medical composition in the form of jelly of the present invention includes ascorbic acid, disodium edetate, tocopherol, etc. The corrective includes citric acid, malic acid, lactic acid, succinic acid, tartaric acid, ascorbic acid, a citrate, a malate, a lactate, a succinate, a tartarate, etc. The sweetening agent includes glucose, fructose, saccharin sodium, sucrose, D-sorbitol, D-mannitol, hydrogenated maltose starch syrup, etc. The emulsifying agent includes polyoxyethylene sorbitan monooleate, sodium lauryl sulfate, etc. The dispersion agent includes an aqueous high molecular weight compound, such as carboxymethylcellulose, sodium alginate, hydroxypropylcellulose, hydroxyethylcellulose, etc. The preservative

includes methyl parahydroxybenzoate (methylparaben), ethyl parahydroxybenzoate (ethylparaben), etc. The flavor includes ones such as menthols, fruit juices, or essential oils. The coloring agent includes caramel, etc.

5           The raw material contained in the Chinese herbal medical composition in the form of jelly of the present invention is not limited as long as it is an ordinary Chinese herbal medicine. For example, Kakkon-to (葛根湯), Sho-seiryu-to (小青竜湯), Sho-saiko-to (小柴胡湯), Hachimi-jio-gan (八味地黄丸), Hochu-ekki-to (補中益氣湯), Sho-kenchu-to (小建中湯), Shofu-san (消風散), Seijo-bofu-to (清上防風湯), Bofu-tsusho-san (防風通聖散), Gorei-san (五苓散), Boi-ogi-to (防已黃耆湯), Otsuji-to (乙字湯), Toki-shakuyaku-san (當歸芍藥散), Keishi-bukuryo-gan (桂枝茯苓丸), Anchu-san (安中散), Heii-san (平胃散), etc., are illustrated. Further, a variety of Chinese herbal medicines are usable as well. The raw material is not only limited to Chinese herbal medicines, but also the raw material made of natural plants is usable in the Chinese herbal medical composition in the form of jelly of the present invention. The raw material selected from Chinese herbal medicines and natural plants is usable in combination with other active substances in the Chinese herbal medical composition in the form of jelly of the present invention. For example, a combination of a cold medicine, an antitussive, an expectorant, and/or a medicine for stomach with a western medicine is usable.

25           The raw material is not limited as long as it is a usual Chinese herbal medicine or natural plant. For example, a crude drug, liquid extract, condensed extract, dry extract, soft extract, fluid extract, etc., are illustrated. The amount of the raw material may be contained in order that the suitable dosage is obtained when a Chinese herbal



medical composition in the form of jelly is taken in the defined amount.

The method for preparing the Chinese herbal medical composition in the form of jelly of the present invention can be the same manner as in a usually known method for jelly preparations. For example, the

5 suitable amount of warmed water as a dispersion medium is added to a base and a raw material and if necessary, a desired substance, and the mixture is stirred to be dispersed, dissolved or suspended, or otherwise the suitable amount of water or cold water as a dispersion medium is added to a base and a raw material and if necessary, a desired  
10 substance at room temperature. The mixture is stirred under warming to be dispersed, dissolved or suspended and the resulting drug is cooled to prepare the Chinese herbal medical composition in the form of jelly. Further, when an ingredient which is not preferable to be exposed to high temperature is contained among a base and a raw material and if  
15 necessary, a desired substance, it may be added after the dispersion, the solution or the suspension prepared above is adjusted to moderate temperature to prepare the Chinese herbal medical composition in the form of jelly. Otherwise an ingredient which is not preferable to be exposed to high temperature may be added thereto just before cooling to  
20 prepare the Chinese herbal medical composition in the form of jelly.

The packed vessel for the Chinese herbal medical composition in the form of jelly of the present invention is not specifically limited, but a stick-like vessel or a bag-like vessel is preferable in respect of carrying and taking it.

25

#### Examples

The present invention is explained by following examples and is not limited by these examples.

#### Examples 1~4 and Comparative examples 1~6

In regard to Examples 1~4 and Comparative examples 1, 5 and 6, the ingredients shown in Tables 1-1 and 1-2 were weighed and each ingredient was dissolved under heating at 80°C. The resulting solution was poured into a stick-like vessel sealed at three parts and cooled to prepare a Chinese herbal medical composition.

In regard to Comparative examples 2~4, after sodium alginate was homogenously dissolved in water, it was warmed at 50~60°C, and thereto were added aqueous dry extract of Kakkon-to (葛根湯). After the mixture was homogeneously dissolved for about 5 minutes, other residual ingredients were added thereto and stirred homogeneously. The mixture was poured into a stick-like vessel sealed at three parts and was cooled to prepare a Chinese herbal medical composition.

The preparation of Example 1 resulted in a good Chinese herbal medical composition in the form of jelly, but the preparation of Comparative example 1 became like a rice cake without forming a jelly. The preparations of Examples 2~4 and Comparative example 5 resulted in Chinese herbal medical compositions in the form of jelly, but the preparations of Comparative examples 2~4 containing sodium alginate did not form jelly. The preparation of Comparative example 6 containing gelatin resulted in a Chinese herbal medical composition in the form of jelly in a refrigerator, but the preparation was a semi-solid at room temperature.

#### Examples 5~7

The ingredients shown in Table 2 were weighed and each ingredient was dissolved under heating at 80°C. The resulting solution was poured into a stick-like vessel sealed at three parts and was cooled

to prepare a Chinese herbal medical composition in the form of jelly.

Examples 8~10 and Comparative examples 7~9

5 In regard to Examples 8~10 and Comparative examples 8 and 9,  
the ingredients shown in Table 3 were weighed and each ingredient was  
dissolved under heating at 80°C. The resulting solution was poured  
into a stick-like vessel sealed at three parts and was cooled to prepare a  
Chinese herbal medical composition in the form of jelly.

10 In regard to Comparative example 7, after sodium alginate was  
homogenously dissolved in water, it was warmed at 50~60°C, and  
thereto were added aqueous dry extract of Seijo-bofu-to (清上防風湯).  
After the mixture was homogeneously dissolved for about 5 minutes,  
other residual ingredients were added thereto and stirred  
homogeneously. The mixture was poured into a stick-like vessel sealed  
15 at three parts and was cooled to prepare a Chinese herbal medical  
composition in the form of jelly.

The preparations of Examples 8~10 resulted in good Chinese  
herbal medical compositions in the form of jelly. The preparations of  
Comparative example 7 and Comparative example 8 containing sodium  
20 alginate and agar, respectively resulted in a Chinese herbal medical  
composition in the form of jelly. The preparation of Comparative  
example 9 containing gelatin resulted in a Chinese herbal medical  
composition in the form of jelly in a refrigerator, but the preparation  
was a semi-solid at room temperature. The preparation of Comparative  
25 example 9 was dissolved in a mouth and gave the bitter taste when it  
was taken, and did not give good feeling (see Table 7). The preparation  
did not serve as a medicine.

### Measurement of the amount of syneresis and the strength of jelly

According to the method below, the Chinese herbal medical compositions in the form of jelly obtained were stored at 40°C under 75 % RH and at 25°C under 60 % RH, respectively. The amount of syneresis and the strength of jelly were measured on each sample and their appearances were observed.

### The method for measuring the amount of syneresis

The method was carried out by standing on end a stick-like packed vessel sealed at three parts into which a Chinese herbal medical composition in the form of jelly was poured (Fig. 1). The ratio of the weight of syneresis remained in the air portion per total amount was calculated.

### The method for measuring the strength on jelly

The method was carried out by taking out a Chinese herbal medical composition in the form of jelly (sample) from a stick-like packed vessel sealed at three parts (see Fig. 1) after it was stored at 25°C for 24 hours, and the sample was measured at 25°C using a tool below.

Measuring tool: Rheometer CR-200D (prepared by San Kagaku)

Measuring conditions: Pressed speed, 30mm/min

Pressure-sensitive axis: Cross section 5 x 40mm x height 15mm  
(Stainless)

### Results

The results obtained on syneresis of the preparations in the form of jelly of Examples 2~4 and Comparative example 5 were shown in

Table 4-1. The preparations of Examples 2~4 were stored both at 40°C under 75 % RH and at 25°C under 60 % RH, and they hardly showed syneresis and their appearances were good. On the other hand, the preparation of Comparative example 5 containing agar showed much  
 5 syneresis both at 40°C under 75 % RH and at 25°C under 60 % RH. Its appearance was bad and did not serve as a medicine.

The results obtained on the jelly strength on the preparations in the form of jelly of Examples 2~4 and the preparation of Comparative example 5 were shown in Table 4-2. Changes on the jelly strength on  
 10 the preparations of Examples 2~4 were not observed under the preservation both at 40°C under 75 % RH and 25°C under 60 % RH. On the contrary, rapid increase of the jelly strength was observed on the preparation Comparative example 5 containing agar under the preservation both at 40°C under 75 % RH and at 25°C under 60 % RH  
 15 in one month.

The preparations of Examples 5~7 resulted in good Chinese herbal medical compositions in the form of jelly, hardly showed syneresis under the preservation conditions both at 0°C under 75 % RH and at 25°C under 60 % RH (see Table 5-1), their appearances were  
 20 good and changes of the jelly strength on them were not observed (see Table 5-2).

The preparations of Examples 8~10 resulted in good Chinese herbal medical compositions in the form of jelly, hardly showed syneresis under the preservation conditions both at 0°C under 75 % RH and at 25°C under 60 % RH (see Table 6-1). Their appearances were  
 25 good and changes of the jelly strength on them were not observed (see Table 6-2).

On the contrary the preparations in the form of jelly of

Comparative examples 7 and 8 showed much syneresis under the preservation conditions both at 0°C under 75 % RH and at 25°C under 60 % RH (see Table 6-1). Their appearances were bad and the jelly strength on them was greatly changed in one month and they could serve as a medicine (see Table 6-2).

As mentioned above, it was ascertained that the Chinese herbal medical composition in the form of jelly of the present invention hardly shows syneresis for a long time and is superior in the preservative stability compared to a jelly preparation containing either gelatin or sodium alginate as a base.

When sodium alginate is used as a base, the Chinese herbal medical composition in the form of jelly is obtained or not obtained depending on the raw material. On the contrast, according to the present invention the good Chinese herbal medical composition in the form of jelly can be obtained regardless of the raw material.

#### Masking effect on bitter taste

By using Chinese herbal medical compositions in the form of jelly and a semi-solid preparation containing gelatin (Comparative example 9), which had the ingredients shown in Table 3, the organoleptic test was carried out. The panel tests were carried out by using 10 persons (5 males and 5 females) and the evaluation were shown as follows:  
++ : too bitter to take one, + : bitter, ± : slightly bitter, - : scarcely bitter, -- : no bitter

A preparation which was prepared by dissolving aqueous dry extract of Seijo-bofu-to (清上防風湯) 7g in water (100g) was used as a control.

As shown in Table 7, in regard to the Control and the

preparations of Comparative examples 8 and 9, almost all persons answered with "too bitter to take them" (Control and Comparative example 9: 10/10, Comparative example 8: 8/10). On the contrast, in regard to the preparation of Example 10, persons who answered with

5 "too bitter to take it" and "bitter" were 3/10 and 7/10, respectively and these preparations showed better result comparing with the preparations of Control and Comparative examples 8 and 9.

The preparation of Comparative example 7 gave the almost same result as the preparation of Example 10, and persons who answered

10 with "too bitter to take it" and "bitter" were 1/10 and 9/10, respectively. The preparation of Comparative example 7 like the preparation of Example 10 showed a better result than the aqueous solution (control), jelly preparations containing agar and gelatin, respectively (Comparative example 8 and 9). Although it is known that the bitter taste is masked

15 by adding sodium alginate to a bitter taste substance, it seems not to exhibit any masking effect as the extract was much.

In regard to the preparation of Example 9, persons who answered with "slightly bitter" were 7/10 and there was none who answered with "too bitter to take it". In regard to the preparation of Example 8,

20 persons who answered with "slightly bitter" were 8/10 and there was none who answered with "too bitter to take it" or "bitter".

As mentioned above, it was ascertained that a Chinese herbal medical composition in the form of jelly wherein the bitter taste of a Chinese herbal medicine effectively masks was obtainable. Especially

25 by solely using  $\kappa$ -carrageenan, it was confirmed that the bitter taste of a Chinese herbal medicine which was contained in the high concentration was excellently masked. Furthermore, by adding a sweetening agent to the Chinese herbal medical composition in the form of jelly prepared by

the present invention, it became possible to take it almost without taking care of the bitter taste of a Chinese herbal medicine.

Table 1-1

Ingredient	Amount (weight %)			
	Ex. 1	Ex. 2	Ex. 3	Ex. 4
Aqueous dry extract of Kakkon-to (葛根湯)	30	15	15	15
ι Carrageenan	0.5	1	1	-
κ Carrageenan	-	-	0.05	1
Carob bean gum	0.1	0.25	0.25	0.25
Xanthan gum	0.2	0.45	0.45	0.45
Sodium alginate	-	-	-	-
Calcium monohydrogen phosphate	-	-	-	-
Glucono-δ-lactone	-	-	-	-
Agar	-	-	-	-
Gelatin	-	-	-	-
Powdered hydrogenated maltose starch syrup	-	6	6	6
D-Sorbitol	-	6	6	6
Glycerin	-	6	6	6
Propylene glycol	-	1	1	1
Propyl parahydroxybenzoate	0.02	0.02	0.02	0.02
Purified water	69.18	64.28	64.23	64.28
Total	100	100	100	100



Table 1-2

Ingredient	Amount (weight %)					
	Comp. ex.1*	Comp. ex.2*	Comp. ex.3*	Comp. ex.4*	Comp. ex.5	Comp. ex.6*
Aqueous dry extract of Kakkon-to (葛根湯)	65	5	15	15	15	15
ι Carrageenan	0.5	-	-	-	-	-
κ Carrageenan	-	-	-	-	-	-
Carob bean gum	0.1	-	-	-	-	-
Xanthan gum	0.2	-	-	-	-	-
Sodium alginate	-	0.8	0.8	2	-	-
Calcium monohydrogen phosphate	-	0.2	0.2	0.5	-	-
Gluclo-δ-lactone	-	2.2	2.2	-	-	-
Agar	-	-	-	-	3	-
Gelatin	-	-	-	-	-	7.5
Powdered hydrogenated maltose starch syrup	-	-	-	-	-	-
D-Sorbitol	-	-	-	-	-	-
Glycerin	-	-	-	-	-	-
Propylene glycol	-	-	-	-	-	-
Propyl parahydroxybenzoate	0.02	0.02	0.02	0.02	0.02	0.02
Purified water	34.18	91.78	81.78	82.48	81.98	77.48
Total	100	100	100	100	100	100

\* : not solidified

Table 2

Ingredient	Amount (weight %)		
	Ex. 5	Ex. 6	Ex. 7
Soft extract of Hachimi-jio-gan (八味地黄丸)	14	-	-
Keishi-bukuryo-gan (桂枝茯苓丸) (crude drug)	-	14	14
ι Carrageenan	1	1	1
κ Carrageenan	-	-	0.1
Carob bean gum	0.2	0.25	0.25
Xanthan gum	0.4	0.45	0.45
Powdered hydrogenated maltose starch syrup	6	6	6
D-Sorbitol	6	6	6
Glycerin	6	6	6
Propylene glycol	1	1	1
Propyl parahydroxybenzoate	0.02	0.02	0.02
Purified water	65.38	65.28	65.18
Total	100	100	100

Table 3

Ingredient	Amount (weight %)							
	Ex. 8	Ex. 9	Ex.10	Comp. ex. 7	Comp. ex. 8	Comp. ex. 9*	Control	
Aqueous dry extract of Seijo-bofu-to (清上防風湯)	14	14	14	14	14	7	7	
ι Carrageenan	1	1	-	-	-	-	-	
κ Carrageenan	-	-	1	-	-	-	-	
Carob bean gum	0.2	0.2	0.2	-	-	-	-	
Xanthan gum	0.4	0.4	0.4	-	-	-	-	
Sodium alginate	-	-	-	2	-	-	-	
Calcium monohydrogen phosphate	-	-	-	0.5	-	-	-	
Agar	-	-	-	-	3	-	-	
Gelatin	-	-	-	-	-	7.5	-	
Powdered hydrogenated maltose starch syrup	6	-	-	-	-	-	-	
D-Sorbitol	6	-	-	-	-	-	-	
Glycerin	6	-	-	-	-	-	-	
Propylene glycol	1	-	-	-	-	-	-	
Propyl parahydroxybenzoate	0.02	0.02	0.02	0.02	0.02	0.02	0.02	
Purified water	65.38	84.38	84.38	83.48	82.98	85.48	92.98	
Total	100	100	100	100	100	100	100	

\* : not solidified

Table 4-1

Amount of syneresis (weight %)	40°C 75 % RH			
	Example 2	Example 3	Example 4	Comp. ex. 5
After 2 days	-	-	-	-
After one month	0.2 %	0.4 %	3.3 %	8.2 %
After 3 months	0.3 %	1.5 %	4.0 %	-
After 6 months	0.8 %	1.9 %	4.4 %	-
Amount of syneresis (weight %)	25°C 60 % RH			
	Example 2	Example 3	Example 4	Comp. ex. 5
After 2 days	no	no	2.3 %	3.0 %
After one month	0.4 %	0.6 %	5.0 %	9.2 %
After 3 months	1.2 %	2.0 %	4.8 %	-
After 6 months	1.0 %	2.3 %	5.4 %	-

Table 4-2

Strength	40°C 75 % RH			
	Example 2	Example 3	Example 4	Comp. ex. 5
Before starting preservation	293g	300g	90g	143g
After one month	290g	314g	88g	191g
After 3 months	302g	322g	84g	-
After 6 months	297g	323g	86g	-
Strength	25°C 60 % RH			
	Example 2	Example 3	Example 4	Comp. ex. 5
Before starting preservation	293g	300g	90g	143g
After one month	285g	304g	87g	180g
After 3 months	297g	308g	88g	-
After 6 months	293g	322g	84g	-

Table 5-1

Amount of syneresis (weight %)	40°C 75 % RH			25°C 60 % RH		
	Ex. 5	Ex. 6	Ex. 7	Ex. 5	Ex. 6	Ex. 7
After 2 days	no	no	no	no	no	no
After one month	no	no	no	0.3 %	no	no
After 3 months	0.3 %	no	no	0.7 %	no	no
After 6 months	0.5 %	no	no	1.1 %	no	no

Table 5-2

Strength	40°C 75 % RH			25°C 60 % RH		
	Ex. 5	Ex. 6	Ex. 7	Ex. 5	Ex. 6	Ex. 7
Before starting preservation	163g	158g	160g	163g	158g	167g
After one month	189g	155g	161g	169g	158g	165g
After 3 months	182g	157g	164g	181g	160g	160g
After 6 months	180g	164g	166g	178g	159g	163g

Table 6-1

Amount of syneresis (weight %)	40°C 75 % RH				
	Ex. 8	Ex. 9	Ex. 10	Comp. ex. 7	Comp. ex. 8
After 2 days	-	-	-	-	-
After one month	0.2 %	0.2 %	4.2 %	9.0 %	10.0 %
After 3 months	0.5 %	0.4 %	5.2 %	-	-
After 6 months	0.8 %	0.5 %	5.0 %	-	-
Amount of syneresis (weight %)	25°C 60 % RH				
	Ex. 8	Ex. 9	Ex. 10	Comp. ex.7	Comp. ex.8
After 2 days	0.2 %	0.2 %	3.3 %	0.9 %	4.0 %
After one month	1.1 %	1.2 %	4.4 %	8.4 %	9.7 %
After 3 months	1.0 %	1.3 %	5.1 %	-	-
After 6 months	1.2 %	1.4 %	5.2 %	-	-

Table 6-2

Strength	40°C 75 % RH				
	Ex. 8	Ex. 9	Ex. 10	Comp. ex. 7	Comp. ex. 8
Before starting preservation	276g	109g	103g	105g	68g
After one month	292g	122g	100g	69g	153g
After 3 months	286g	118g	111g	-	-
After 6 months	-	296g	114g	-	-
Strength	25°C 60 % RH				
	Ex. 8	Ex. 9	Ex. 10	Comp. ex.7	Comp. ex.8
Before starting preservation	276g	109g	103g	105g	68g
After one month	287g	111g	114g	64g	139g
After 3 months	284g	110g	116g	-	-
After 6 months	282g	118g	118g	-	-

Table 7

Olganoleptic test	Evaluation (Unit: person)				
	++	+	±	-	--
Ex. 8	0	0	2	8	0
Ex. 9	0	3	7	0	0
Ex. 10	3	7	0	0	0
Comp. ex. 7	1	9	0	0	0
Comp. ex. 8	8	2	0	0	0
Comp. ex. 9	10	0	0	0	0
Control	10	0	0	0	0

++ : too bitter to take one

+ : bitter

± : slightly bitter

- : scarcely bitter

-- : no bitter

#### INDUSTRIAL APPLICABILITY

In the present invention, by using at least one substance selected from the group consisting of carrageenan, carob bean gum and xanthan gum as a base (not containing phosphate buffer) of the jelly preparation containing a Chinese herbal medicine, a Chinese herbal medical composition in the form of jelly, which hardly causes syneresis, is superior in the preservative stability, is broadly applicable to a Chinese herbal medicine, and is orally taken without taking care of bitter, etc., of a Chinese herbal medicine is obtainable. Furthermore, even when the Chinese herbal medical composition in the form of jelly of the present invention can stand for the test for medicines, for example the long term-preservation test at 25°C under 60 % RH for 3 years, and the accelerated preservation test at 40°C under 75 % RH for 6 months and

is guaranteed in the preservative stability.